

Pain Control

**+: AEMT dose =
half max**

- B**
- Ensure pulse oximetry
 - As appropriate, begin non-analgesic comfort measures:
 - Splinting
 - Blankets
 - Temperature control
- B**

ADULT

A IV/IO Access **A**

PEDIATRIC

A

Acetaminophen
THEN
Ketorolac
OR
Morphine
OR
Fentanyl
BEFORE
Ketamine

A

Consider:
Ondansetron

P

Consider:
Promethazine

P

**AEMT max dosing is half
of overall max dose. When
exceeded, patient care
transfers to the Paramedic.**

A

Acetaminophen
BEFORE
Morphine
BEFORE
Fentanyl

A

Consider:
Ondansetron

P

Ketorolac

P

ADULT:

Acetaminophen
500 mg
Fentanyl +
50-100 mcg every 10 min, Max 200 mcg
Morphine +
2-4 mg every 10 min, Max 8 mg
Ketamine
0.1-0.3 mg/kg IV/IO every 15 min, Max 100 mg
50-100 mg IN
Ketorolac
30 mg IV | 60 mg IM
Ondansetron
4 mg, Max 8 mg
Promethazine
12.5-25 mg IM, Max 25 mg

PEDIATRIC:

Acetaminophen
15 mg/kg
Fentanyl +
1 mcg/kg every 10 min, Max 100 mcg
Morphine +
0.1 mg/kg single dose, Max 4 mg
Ketorolac
> 2 yrs old: 0.5 mg/kg, Max 30 mg
Ondansetron
0.1 mg/kg, Max 4 mg